

A NOTE ON ARTIN'S CONSTANT

IVAN CHEREDNIK [†]

We suggest a representation of Artin's constant, which conjecturally describes the density of prime p such that “generic” $g \in \mathbb{Z}$ is primitive modulo p . Namely, $A = \lim_{N \rightarrow \infty} R_k(N)$ for $R_k(N) = \frac{\sum p^k \phi(p_i - 1)}{\sum p^k (p_i - 1)}$, where the summation is over first N prime p_i , $k \in \mathbb{Z}_+$. The classical summation formula is as follows: $A = \lim_{N \rightarrow \infty} \Sigma(N)$, where $\Sigma(N) = \frac{1}{N} \sum \frac{\phi(p_i - 1)}{p_i - 1}$. The changes needed for arbitrary g are addressed in Theorem 1, a good exercise in basic analytic and algebraic number theory. The same procedure can be applied to other number-theoretic constants like A (see, e.g., [Ni]). In Theorem 2, we demonstrate how it works for the Stephens constant and for Artin's constants of higher ranks (for the density of prime p such that a given set of “generic” integers generates $\mathbb{Z}_p^*$).

The following three features of this approach vs. the summation formulas are worth noticing.

1) The *restricted summation* suggested by P. Moore to make the Σ -formula matching the right heuristic density for arbitrary $g \in \mathbb{Z}$ gives the desired answer in our approach only when g is not a pure (odd) power in $\mathbb{Z}$. Otherwise, nontrivial rational multiplicative corrections occur; they are calculated in Theorem 1, (ii).

2) The p^k -terms in the denominator and numerator of $R_k(N)$ do not influence the limit, which can be heuristically associated with switching from primitive roots in $\mathbb{Z}_p$ to those in $(\mathbb{Z}/(p^{k+1}))^*$. The extra p^k -factors disappear (cancel) in the corresponding summation formula. When $k = 0$, our R -formula for A (without restricting the summation) follows from [Pi].

3) The R -formulas oscillate significantly around A (and the other constants). The magnitude of *oscillations* increases as k grows; see Figure 1. Representing $R_k(N) = \sum_{i=1}^N w_i \frac{\phi(p_i - 1)}{p_i - 1}$, the weights w_i change from $O(\frac{\log N}{N^{k+2}})$ for small i to $O(\frac{(\log N)^{k+1}}{N})$ for $i \sim N$. Thus, large p

[†] March 11, 2009 Partially supported by NSF grant DMS-0800642.

receive greater weights in our approach, especially when k is large, which increases the range of oscillations.

1. Brief history. Artin's primitive root conjecture states that given an integer g , possibly negative but not a perfect square in $\mathbb{Z}$, the number $\mathcal{P}_N(g)$ of prime p among $p_1 = 2, p_2 = 3, \dots, p_N$ such that g is primitive modulo p approaches asymptotically $A(g)N$ as $N \rightarrow \infty$ for

$$(1) \quad A(g) = A_h A_d, \quad A_d = \left(1 - \mu(|d|) \prod_{p|d, p|h} \frac{1}{p-2} \prod_{p|d, p \nmid h} \frac{1}{p^2 - p - 1}\right),$$

$$A_h = \prod_{p|h} \left(1 - \frac{1}{p-1}\right) \prod_{p \nmid h} \left(1 - \frac{1}{p(p-1)}\right), \quad \text{where}$$

$d = \text{Discriminant}(\mathbb{Q}[\sqrt{g}])$, $g = g_o^h$ for $g_o \in \mathbb{Z}$ and maximal $h \in \mathbb{N}$.

Note that $\mu(|d|) = 0$ and, respectively, $A(g) = A_h$ if and only if the discriminant d is not from $1 + 4\mathbb{Z}$. If $h = 1$ for such g , then $A(g)$ equals

$$(2) \quad \text{Artin's constant} = A = \prod_{\text{prime } p} \left(1 - \frac{1}{p(p-1)}\right).$$

According to [St], Artin's conjecture was finalized around 1965. In 1967, it was deduced by Hooley [Ho] from the generalized Riemann hypothesis for the fields $K_m = \mathbb{Q}[\zeta_m, g^{1/m}]$ for squarefree m . See [Mo1] for a comprehensive introduction (including some recent developments). See also [Mu, Le].

Artin's heuristic approach to this conjecture was based on the expectation that events “*prime p does not split completely in K_q for prime q* ” are independent (subject to later qualitative and quantitative corrections). For instance, one can expect that A equals $\lim_{N \rightarrow \infty} \mathcal{P}_N(g)/N$ if g is *generic* as far as primitive roots modulo prime p are concerned. It leads to the following heuristic summation formula for Artin's constant:

$$(3) \quad A = \lim_{N \rightarrow \infty} \Sigma(N), \quad \Sigma(N) = \frac{1}{N} \sum_{i=1}^N \frac{\phi(p_i - 1)}{p_i - 1},$$

which can be checked unconditionally, without any reference to Artin's conjecture. See [LL], [Mo1]. P. Moore extended it to arbitrary $A(g)$ by switching to p in this summation such that g is a quadratic nonresidue modulo p and $(p-1, h) = 1$ (heuristically, it makes sense); see below.

As Lehmers wrote, the convergence in (3) is “discouragingly slow” (they considered prime numbers $p < 1500000$). It remains very slow when prime numbers in much greater ranges are considered, generally, no better than the (conjectural) convergence of $\mathcal{P}_N(2)/N$ to $A = A(2)$; cf. Table “Artin’s constant estimates” from [Si] ($p < 10^{14}$).

2. Main Theorem. A refine version of this heuristic approach is from [Mo1] (for any integer g):

$$(4) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \varphi_g(p_i) = A(g) \quad (\stackrel{\text{RH}}{=} \lim_{N \rightarrow \infty} \mathcal{P}_N(g)/N), \quad \text{where}$$

$$\varphi_g(p) \stackrel{\text{def}}{=} 2 \frac{\phi(p-1)}{p-1} \quad \text{for} \quad \left(\frac{g}{p}\right) = -1 \quad \text{and} \quad (p-1, h) = 1,$$

$$\varphi_g(p) \stackrel{\text{def}}{=} 0 \quad \text{otherwise}.$$

Heuristically, it is equally reasonable to expect that

$$(5) \quad A = \lim_{N \rightarrow \infty} \frac{\sum_{i=1}^N \phi(p_i - 1)}{\sum_{i=1}^N (p_i - 1)}$$

for sufficiently general g . Switching here to the restricted summation from (4), we come to the following theorem.

Theorem 1. (i) For arbitrary integers $k \geq 0$ and g (possibly, negative),

$$(6) \quad A(g) = \lim_{N \rightarrow \infty} \frac{\sum_{i=1}^N p_i^k \phi(p_i - 1)}{\sum_{i=1}^N p_i^k (p_i - 1)},$$

provided that $h = 1$, where the summation Σ' is over prime p_i such that $\left(\frac{g}{p_i}\right) = -1$ and $(p_i - 1, h) = 1$.

(ii) If $h > 1$, then the limit in the r.h.s. of (6) equals

$$(7) \quad \tilde{A}(g) = \tilde{A}_h \tilde{A}_d, \quad \text{where} \quad \tilde{A}_h = \prod_{p \nmid h} \left(1 - \frac{1}{p(p-1)}\right),$$

$$\tilde{A}_d = 1 \quad \text{if} \quad d \mid h \quad \text{and} \quad \tilde{A}_d = A_d \quad \text{otherwise}.$$

Sketch of proof. We follow [LL, Mo1], restricting ourselves with (5); see also [Pi]. Coupling the generalized Landau formula from [SZ]

$$(8) \quad \sum_{p \leq x} p^m = \frac{(1 + o(1)) x^{m+1}}{(1 + m) \log x}, \quad m \geq 0,$$

with the classical estimate

$$(9) \quad \frac{\pi(x, d, 1)}{\pi(x)} = \frac{1 + O(1/\log x)}{\phi(d)},$$

where $\pi(x, d, 1)$ is the number of prime numbers $p \leq x$ in $1 + d\mathbb{N}$ ($\pi(x) = \pi(x, 1, 1)$), one arrives at:

$$(10) \quad \frac{\pi^{(m)}(x, d, 1)}{\pi^{(m)}(x)} = \frac{1 + O(x^m/\log x)}{\phi(d)},$$

where $\pi^{(m)}(x, d, 1) = \sum_{p \leq x} p^m$ over prime $p \in 1 + d\mathbb{N}$. Then,

$$\begin{aligned} \sum_{p \leq x} \phi(p-1) &= \sum_{p \leq x} \sum_{d|p-1} (p-1) \frac{\mu(d)}{d} \\ &= \sum_{d|p-1} \frac{\mu(d)}{d} \sum_{p \leq x} (p-1) = \sum_{d \leq x} \frac{\mu(d)}{d} (\pi^{(1)}(x, d, 1) - \pi(x, d, 1)). \end{aligned}$$

Finally,

$$\frac{\sum_{p \leq x} \phi(p-1)}{\sum_{p \leq x} (p-1)} \sim \sum_{d \leq x} \frac{\mu(d)}{d\phi(d)} \xrightarrow{x \rightarrow \infty} A.$$

We will omit the arguments (from the basic algebraic number theory) that give the rational corrections for arbitrary g . $\square$

In (5,6), only the leading powers of p_i matter; for instance, one can take here the ratio $\sum_p \phi(\phi(p^{k+1}))/\sum_p \phi(p^{k+1})$, the heuristic probability for g being a primitive root modulo p^{k+1} over all prime p , which leads to the same Artin constant.

3. Further examples. Let us apply the same procedure to Stephens' constant $S(a, b)$ and the higher rank Artin constants. The former is defined for a given pair $a, b \in \mathbb{Q}^*$ such that $a^r b^s = 1 \implies r = 0, s = 0$ for $r, s \in \mathbb{Z}$; it describes the density of prime p such that $b = a^m \pmod p$ for some $m \in \mathbb{Z}$. Modulo the generalized Riemann hypothesis, it equals

$$(11) \quad C_{ab} S \quad \text{for} \quad S \stackrel{\text{def}}{=} \prod_{i=1}^{\infty} \left(1 - \frac{p_i}{p_i^3 - 1}\right),$$

where the factors C_{ab} are rational [S, MS]. These factors were calculated explicitly in [MS] under the condition that the group $\mathbb{Q}^*/\langle a, b, -1 \rangle$

is torsion free. Assuming that a, b are “random”, the heuristic probability $\mathcal{BS}(p)$ that $b = a^m \pmod{p}$ for some m can be readily calculated:

$$(12) \quad \mathcal{BS}(p) = \mathcal{S}(p)/(p-1)^2, \text{ where } \mathcal{S}(p) \stackrel{\text{def}}{=} \sum_{d|p-1} d\phi(d) \\ = \prod_{j=1}^m \frac{q_j^{2k_j+1} + 1}{q_j + 1} \text{ for prime factorization } p-1 = \prod_{j=1}^m q_j^{k_j}.$$

We naturally omit the prime numbers that divide the numerators or denominators of a, b .

The rank r Artin constant $A_r(g_1, \dots, g_r)$ describes the heuristic density of prime p such that a given set of nonzero integers $\{g_1, \dots, g_r\}$ (or rationals) generates $\mathbb{Z}_p^*$. See [CP]. Its “generic” value (modulo the generalized Riemann hypothesis) is as follows:

$$(13) \quad A_r \stackrel{\text{def}}{=} \prod_{i=1}^{\infty} \left(1 - \frac{1}{p_i^r (p_i - 1)} \right).$$

The corresponding probability at p for “random” $g_1, \dots, g_r$ equals

$$(14) \quad PA_r(p) = \mathcal{A}_r(p)/(p-1)^r \text{ for } \mathcal{A}_r(p) \stackrel{\text{def}}{=} \prod_{j=1}^m (q_j^{rk_j} - q_j^{r(k_j-1)})$$

in terms of the prime factorization $p-1 = \prod_{j=1}^m q_j^{k_j}$.

The summation (unconditional) limiting formulas are as follows:

$$(15) \quad S = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \mathcal{BS}(p_i), \quad A_r = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N PA_r(p_i).$$

The proof of the following theorem is similar to that of Theorem 1.

Theorem 2. *For an arbitrary integer $k \geq 0$,*

$$(16) \quad S = \lim_{N \rightarrow \infty} \frac{\sum_{i=1}^N p_i^k \mathcal{S}(p_i)}{\sum_{i=1}^N p_i^k (p_i - 1)^2},$$

$$(17) \quad A_r = \lim_{N \rightarrow \infty} \frac{\sum_{i=1}^N p_i^k \mathcal{A}_r(p_i)}{\sum_{i=1}^N p_i^k (p_i - 1)^r},$$

where the summation is over consecutive prime numbers p_i . □

4. Numerical aspects. For the constants C considered above, we plot $\frac{\Sigma(N)}{C} - 1$, shown blue-thin, for the classical summation $\Sigma(N)$ and $\frac{R(N)}{C} - 1$, which are red-thick, for our ratio approximations $R(N)$; for instance, $\Sigma(N)$ is from (3) for Artin's A . The range is $N \leq 1000M$.




FIGURE 1. Deviation from A as $k = 0, 1$ for 1000M primes.

Figure 1 compares the stabilization of (3), blue-thin, to the Artin constant and the stabilization of (5), red-thick, as $k = 0, 1$. The function $\Sigma(N) - A$ remains positive in the range $N \leq 1000M$; $R_k(N)$ oscillate around Artin's constant $A \approx 0.37395581361920228805$ (the zero level of this graph). The amplitude of oscillations become larger for $R_{k=1}$ vs. $R_{k=0}$, however the graphs are very much similar.

We note that the best way to calculate A and similar constants is based on the known product formulas in terms of $\zeta(n)$ for integers $n > 1$.






FIGURE 2. Deviation from $A(5)$ for 1000M primes.

Figure 2 shows the convergence of (4) and (6) to $A(g = 5)$ for $k = 0$; notice that the blue-thin curve remains beyond the red-thick one in this range.

The last two plots show the graphs for the Stephen constant $S \approx 0.57595996889294543964$, Figure 3, and the rank 3 Artin constant $A_3 \approx 0.85654044485354217443$, Figure 4; here $k = 0$. The convergence rate and other features of these four graphs are similar to those for A and $A(5)$. There is striking (qualitative) similarity of these two figures, including the oscillations, although the convergence rate in Figure 4 (for A_3) is significantly (almost 10 times) greater than in Figure 3.

DEVIATION FROM S , STEP = 5000000 PRIMESFIGURE 3. Deviation from S for 1000M primes.DEVIATION FROM $A[r=3]$, STEP = 5000000 PRIMESFIGURE 4. Deviation from A_3 (rank=3) for 1000M primes.

Qualitatively, the behavior of $R(N)$ for large N can be evaluated following [Pi]. For instance, the functions $|R(N)/A - 1|$ in Figure 1 must be no greater than $C_m(\log N)^{-m}$ for *any* fixed $m > 0$ and proper constant C_m (depending on k) as $N \gg 0$. Generally speaking, C_m can be estimated in terms of (the order of) N , but we will not discuss it. Indeed, the graphs of $R(N)/A - 1$, red-thick from Figure 1, look like $O((\log N)^{-4})$ in the range $N < 1000M$.

The nature of oscillations of the functions $R(N)$ around the corresponding constants remains unclear.

Acknowledgements. I am grateful to Zeev Rudnik for valuable comments. I am very thankful to Pieter Moree for reading the note, suggesting interesting questions toward comparing our approach with the results from [Mo2, Mo3] and for the reference to [Pi].

REFERENCES

- [CP] L. Cangelmi, and F. Pappalardi, *On the r -rank Artin conjecture II*, J. Number Theory 75 (1999), 120–132. [5](#)
- [Ho] C. Hooley, *Artins conjecture for primitive roots*, J. Reine Angew. Math. 225 (1967), 209–220. [2](#)
- [LL] D.H. Lehmer, and E. Lehmer, *Heuristics, anyone ?*, in “Studies in Mathematical Analysis and Related Topics”, 202–210, Stanford Univ. Press, Stanford, CA, 1962. [2](#), [3](#)
- [Le] H.W. Lenstra, jr., *On Artins conjecture and Euclids algorithm in global fields*, Invent. Math. 42 (1977), 202–224. [2](#)
- [Mo1] P. Moree, *Artins primitive root conjecture*, Preprint arXiv:math/0412262 [math.NT], 2004. [2](#), [3](#)
- [Mo2] P. Moree, *Asymptotically exact heuristics for (near) primitive roots*, J. Number Theory 83:1 (2000), 155–181. [9](#)
- [Mo3] P. Moree, *Asymptotically exact heuristics for (near) primitive roots. II*, Japan. J. Math. (N.S.) 29:2 (2003), 143–157. [9](#)
- [MS] P. Moree, and P. Stevenhagen, *A two variable Artin conjecture*, J. Number Theory 85 (2000), 291–304. [4](#)
- [Mu] R. Murty, *Artins conjecture for primitive roots*, Math. Intelligencer 10:4 (1988), 59–67. [2](#)
- [Ni] G. Niklasch, *Some number-theoretical constants arising as products of rational functions of p over primes*, Preprint 2002;
<http://www.gn-50uma.de/alula/essays/Moree/Moree.en.shtml> . [1](#)

- [Pi] S.S. Pillai, On the sum function connected with primitive roots, Proc. Indian Acad. Sci., Sect. A. 13 (1941), 526–529. [1](#), [3](#), [9](#)
- [Si] T. Oliveira e Silva, *Least primitive root of prime numbers*, Preprint 2004; <http://www.ieeta.pt/~tos/p-roots.html> . [3](#)
- [SZ] T. Salát, and S. Znám, *On the sums of prime powers*, Acta Fac. Rer. Univ. Com. Math. 21 (1968), 21–25. [3](#)
- [S] P.J. Stephens, *Prime divisors of second order linear recurrences*, J. Number Theory 8:3 (1976), 313–345. [4](#)
- [St] P. Stevenhagen, *The correction factor in Artins primitive root conjecture*, J. Théor. Nombres Bordeaux 15 (2003), 383–391. [2](#)

(I. Cherednik) DEPARTMENT OF MATHEMATICS, UNC CHAPEL HILL, NORTH CAROLINA 27599, USA, CHERED@EMAIL.UNC.EDU